



Pür Combo is a liquid nutrient well-balanced in magnesium and readily assimilable micronutrients to prevent and treat deficiencies. Pür Combo contains amino acids obtained by enzymatic hydrolysis that stimulate the physiological processes of the plant, absorption and mobility of nutrients in the plant as well as establishing resistance mechanisms against abiotic stresses (drought, low temperature, salinity, etc.). The synergy between the elements activates many vital processes while enhancing the crop's yield inducing factors (cellular division, photosynthesis, flowering, sugar synthesis and transport, hormone and protein synthesis, etc.). Pür Combo must be applied based on soil or tissue analyses or as soon as deficiency symptoms appear. For more information on Pür Combo, consult your agronomic adviser.

agronomic benefits


- Prevents and rectifies many deficiencies
- Broad spectrum of effectiveness on every crop and every type of soil
- Better use of nutritional elements
- Resistance against stresses
- Yield optimization

density 1.261 kg/L

Guaranteed minimum analysis

Magnesium (Mg)	1.2%
Sulphur (S) (actual)	2.5%
Boron (B) (actual)	0.6%
Copper (Cu) (actual)	0.25%

0-0-0 + 2.5% S + micros



Also contains non-plant food ingredients

Amino acids	2.5%
Citric acid	5%

Rate US gallon/acre			Foliar application
Rate L/ha]	
Crops			Application timing and remarks
All	2-4	0.2-0.4	During periods of rapid growth. Apply before deficiency symptoms appear. Water volume: at least 200 L/ha (21.5 gallons/acre).
Fruits trees	2-4	0.2-0.4	As soon as leafage coverage is sufficient. Water volume: at least 600 L/ha (65 gallons/acre). Do not use this product when fruit coloration and maturity may be delayed by nitrogen application.

Liquid nutrient

with micronutrients

Soil application

Consult your agronomic adviser for the appropriate dosage for your conditions.

Apply at suggested rates. Water volume: at least 200 L/ha (21.5 gallons/acre).

directions for use

Mixing and application. Shake well before using. Fill the tank three quarters full. Start agitation and add product. Clean the empty containers and add the rinsing water to the tank. Add the rest of the water and apply without delay. Do not apply under drying conditions and when outside temperature is too high otherwise culture may be damaged.

Tank mixing. In some cases, tank mixing with pest control products can result in physical and chemical incompatibilities which may impact the efficacy of the products and the crop performance. Always read and follow label directions of the tank mix partners. Consult the pesticides manufacturer representatives and an Agro-100 representative to help determine the safety and agronomic outcome of the mix.

Compatibility. This product is compatible with most agrochemical products except phosphates. Check compatibility before mixing in application tank or check with your adviser. **Check compatibility as follow**. In a clean, transparent jar, add the water and other products, one by one in the same proportion as proposed application. Agitate gently, Let the jar sit for 30 minutes. Incompatible products will form flakes, sludge, gels or other precipitates. Separation or layering may appear. Do not apply in case of incompatibility.

Storage. Avoid freezing.

Caution. This product contains boron, copper, iron, manganese, molybdenum and zinc and must be used only as recommended. It may prove harmful when misused. This product must be used as complement to a soil fertilization program based on soil or plant tissue analyses.

Guarantee. The vendor's liability will be limited to the terms of this label. The buyer will then be responsible for all risks and damages to persons or properties resulting from the use of this product, and hereby accepts these conditions.

